

SUPPLEMENTARY MATERIALS FOR THE ARTICLE:

**Habitat associations of the Brown Shrike (*Lanius cristatus* Linnaeus, 1758) during the non-breeding period in human-modified landscapes of Southeast Asia**

Artur Golawski<sup>1\*</sup>, Cezary Mitrus<sup>2</sup>, Przemysław Obłóza<sup>1</sup>, Santi Xayyasith<sup>3</sup>, Zbigniew Kasprzykowski<sup>1</sup>

Folia Biologica (Kraków), vol. 74 (2026), No 1.

DOI: [https://doi.org/10.3409/fb\\_74-1.02](https://doi.org/10.3409/fb_74-1.02)

**SM.02. Supplementary materials\_Fig. S1. Univariate\_plots.**

<sup>1</sup>University of Siedlce, Faculty of Sciences, Prusa 14, 08-110 Siedlce, Poland

<sup>2</sup>Department of Biology and Ecology of Animals, Institute of Biology, University of Environmental and Life Sciences, Chełmońskiego 38c, 51-631 Wrocław, Poland

<sup>3</sup>Faculty of Environmental Sciences, National University of Laos, Vientiane 01170, Lao PDR

\*Corresponding author; e-mail: [artur.golawski@uws.edu.pl](mailto:artur.golawski@uws.edu.pl)




Fig. S1. Exploratory univariate plots showing the raw relationships between Brown Shrike presence and each of the ten habitat variables prior to GLMM modelling.